

**BLACK HOLES AS P-BRANES\***

BENJAMIN C. HARMS

and

YVAN LEBLANC

*Department of Physics and Astronomy, The University of Alabama  
Tuscaloosa, AL 35487-0324, USA***ABSTRACT**

We review briefly the thermodynamical interpretation of black hole physics and discuss the problems and inconsistencies in this approach. We provide an alternative interpretation of black holes as quantum objects and investigate the statistical mechanics of a gas of such objects in the microcanonical ensemble. We argue that the theory of black holes has the conformal properties of duality and satisfaction of the statistical bootstrap condition. We show in the context of mean field theory that the thermal vacuum is the false vacuum for a black hole and define a microcanonical vacuum which leads to a number density characteristic of pure states for the Hawking radiation.

**1. Introduction**

Nearly twenty years ago, Hawking proposed that the laws of quantum mechanics do not hold in the creation and subsequent evaporation of black holes<sup>1</sup>. In this picture, which is based on the premise that black holes can be treated as thermodynamical systems<sup>2</sup>, black holes can radiate by the process of particles tunneling quantum mechanically through the horizon. The laws of quantum mechanics are violated because the emerging radiation always has a number density function which is characteristic of mixed states, while the accreted radiation may have been in pure states<sup>3</sup>. This violates the unitarity principle. During the period since Hawking's original proposal, many papers have been written on the so-called information loss paradox which occurs in the thermodynamical interpretation of black holes<sup>\*</sup>.

In a series of papers<sup>5,6,7,8,9,10,11</sup> we have pointed some inconsistencies in the thermodynamical interpretation of processes involving black holes and have offered an alternative description of black holes. In our description black holes are considered to be quantum objects, specifically, extended quantum objects or p-branes. The main purpose of the present work is to summarize our results to date and to suggest possible lines of research which follow from these results.

---

<sup>\*</sup> To appear in the Proceedings of the 3rd Workshop on Thermal Field Theories and their Applications, Banff, Alberta, Canada, August 15-27, 1993

<sup>\*</sup>For a review of the subject see, for example, Ref.[4]

We begin in section **2.** with a brief summary of the issues in the thermodynamical interpretation of processes involving black holes. We discuss the inconsistencies alluded to above in detail. In section **3.** we discuss our interpretation of the WKB formula as the quantum tunneling probability and review our results for the statistical mechanics of a gas of black holes. In section **4.** we discuss the thermodynamical interpretation of black holes within the context of mean field theory and prove that the thermal vacuum is the false vacuum for a black hole system. In section **5.** we present an alternative vacuum for such a system and prove that the particle number density for the radiated particles derived from this formulation represents a pure state. In the final section we discuss our conclusions and future extensions of our work.

## 2. Thermodynamical Interpretation of Black Holes

Bekenstein<sup>2</sup> was the first to suggest that the area of a 4-dimensional black hole can be identified with its entropy and the surface gravity with its temperature. The area of a classical black hole increases when matter falls into the black hole. Thus as a classical black hole accretes matter, its entropy increases.

Bardeen, Carter and Hawking<sup>12</sup> derived a relation between the mass difference of neighboring equilibrium states of a black hole and the change in its area

$$\Delta M = \kappa \Delta A , \quad (1)$$

and showed that the temperature is related to the surface gravity by

$$T = \frac{\kappa}{2\pi} . \quad (2)$$

By taking quantum effects into account, Hawking<sup>13</sup>, using an operator formalism and Gibbons and Hawking<sup>14</sup>, using the WKB approximation, demonstrated that black holes can radiate. In the WKB approximation a conical singularity develops in the Euclidean spacetime and is removed by requiring that the imaginary time variable be periodic with period  $8\pi M$ ,  $M$  being the mass of the black hole. This value of the imaginary time is identified with the inverse temperature  $\beta_H$ . The partition function for the black hole is given by

$$\begin{aligned} Z &= \text{Tr} e^{-\beta H} \\ &\sim e^{-S_E} , \end{aligned} \quad (3)$$

where  $S_E$  is the classical Euclidean action. The Hawking entropy is given by

$$\begin{aligned} S_H &= \beta M - S_E \\ &= S_E ; \quad (D = 4) . \end{aligned} \quad (4)$$

A calculation of the canonical (inverse) temperature  $\beta$  shows that

$$\beta = \frac{\partial S_H}{\partial E} = \beta_H = 8\pi M . \quad (5)$$

The Hawking entropy is determined from the Euclidean spacetime metric. It is related to the area of the black hole by,

$$S_H = \frac{A}{4} . \quad (6)$$

For a 4-dimensional Schwarzschild black hole the radius of the horizon is  $r_+ = 2M$ , so that

$$S_H(M) = 4\pi M^2 = \frac{1}{16\pi} \beta_H^2 . \quad (7)$$

This interpretation of black holes has severe problems. The first problem is that the canonical specific heat, an intrinsically positive quantity, turns out to be negative

$$C = \frac{\partial E}{\partial T} = -\frac{\beta^2}{8\pi} \quad (8)$$

A second problem is that the partition function as calculated the microcanonical density of states

$$Z = \int e^{-\beta E} \Omega_H dE , \quad (9)$$

where

$$\Omega_H = e^{S_H} , \quad (10)$$

blows up for all temperatures,

$$Z = \int_0^\infty dE e^{-\beta E + 4\pi E^2} \rightarrow \infty , \quad (11)$$

indicating a breakdown of the WKB approximation and therefore the inequivalence of the canonical and microcanonical ensembles for black holes. A third problem is of a quantum mechanical nature. The radiation coming out of the black hole has been shown to have a Planckian distribution

$$n = \frac{1}{e^{M\beta} - 1} . \quad (12)$$

This implies a loss of coherence, because pure states may come into the black hole, but only mixed states come out. This is a statement of the so-called information loss paradox. In the thermodynamical interpretation unitarity is lost and with it one of the basic principles of quantum mechanics. This interpretation requires the abandonment of quantum mechanics and the postulation of new physical laws. The nature of these new laws is unknown at present.

### 3. Quantum Interpretation of Black Holes

The problems with the thermodynamical interpretation of black holes can be avoided by adopting a different interpretation of the WKB formula. If the WKB formula is interpreted to be, as is usual, the tunneling probability per unit volume for a particle to tunnel through the horizon of a black hole

$$P \propto e^{-S_E(M)} , \quad (13)$$

then the black hole can be viewed as the quantum excitation mode of a  $\frac{D-2}{D-4}$ -brane. To see this, one notes that the quantum degeneracy of states for such an object is essentially the inverse of the tunneling probability

$$\sigma(M) \simeq ce^{S_E(M)} , \quad (14)$$

where  $c$  is a constant which is determined by quantum field theoretical corrections.

As an example we consider the Schwarzschild black hole, which in  $D$ -dimensions has a Euclidean metric given by

$$ds^2 = e^{2\Phi} d\tau^2 + e^{-2\Phi} dr^2 + r^2 d\Omega_{D-2}^2 , \quad (15)$$

where

$$e^{2\Phi} = 1 - \left(\frac{r_+}{r}\right)^{D-3} . \quad (16)$$

The Euclidean action calculated from this metric is

$$S_E = \frac{A_{D-2}}{16\pi} \beta_H r_+^{D-3} , \quad (17)$$

with

$$\begin{aligned} \beta_H &= \frac{2\pi}{[e^\Phi \partial_r e^\Phi]_{r=r_+}} = \frac{4\pi r_+}{D-3} \\ M &= \frac{D-2}{16\pi} A_{D-2} r_+^{D-2} , \end{aligned} \quad (18)$$

where  $A_{D-2}$  is the area of a unit  $D-2$  sphere. Eliminating the horizon radius  $r_+$  in favor of the mass, the Euclidean action becomes

$$S_E = C(D) M^{\frac{D-2}{D-3}} , \quad (19)$$

where  $C(D)$  is the dimension-dependent constant

$$C(D) = \frac{4^{\frac{D-1}{D-3}} \pi^{\frac{D-2}{D-3}}}{(D-3)(D-2)^{\frac{D-2}{D-3}} A_{D-2}^{\frac{1}{D-3}}} . \quad (20)$$

Substituting in for  $S_E$  in the quantum degeneracy of states expression we find

$$\sigma(M) \simeq ce^{C(D)M^{D-2/D-3}} . \quad (21)$$

Comparing this expression to those known for nonlocal field theories<sup>15,16,17,18</sup>, we find that it corresponds to the degeneracy of states for an extended quantum object ( $p$ -brane) of dimension  $p = \frac{D-2}{D-4}$ .

Returning to the thermodynamical interpretation, we find for the statistical mechanical density of states

$$\Omega_H(M) = e^{S_H(M)} \quad (22)$$

where  $S_H$  is the Hawking entropy

$$S_H = \beta_H M - \beta_H F(\beta_H) = \beta_H M - S_E . \quad (23)$$

In  $D$ -dimensions we find from Eq.(18) and Eq.(19) that the entropy is

$$S_H = (D-3)S_E(M) , \quad (24)$$

giving for the density of states

$$\Omega_H(M) \simeq \sigma^{D-3}(M) . \quad (25)$$

The partition function obtained from this density of states is

$$\begin{aligned} Z(\beta) &= \int_0^\infty dE e^{-\beta E} e^{(D-3)C(D)E^{\frac{D-2}{D-3}}} \\ &\rightarrow \infty \text{ for } D \geq 4 , \end{aligned} \quad (26)$$

illustrating for  $D$ -dimensional black holes the earlier results for 4-dimensional black holes. The statistical mechanics of black holes must therefore be studied in the more fundamental microcanonical ensemble.

Within the context of our interpretation of black holes as quantum objects we have considered a gas of such objects. The microcanonical ensemble in 4 dimensions is determined by the microcanonical densities

$$\Omega(E, V) = \sum_{n=1}^{\infty} \Omega_n(E, V) , \quad (27)$$

with

$$\begin{aligned} \Omega_n(E, V) &= \left[ \frac{V}{(2\pi)^3} \right]^n \frac{1}{n!} \prod_{i=1}^n \left( \int_{m_0}^{\infty} dm_i \sigma_{BH}(m_i) \int_{-\infty}^{\infty} d^3 p_i \right) \\ &\times \delta \left( E - \sum_{i=1}^n E_i \right) \delta^3 \left( \sum_{i=1}^n p_i \right) , \end{aligned} \quad (28)$$

where  $\sigma_{BH}(m_i)$  is given by Eq.(21) and  $m_0$  is the mass of an extreme black hole ( $m_0 = 0$  for Schwarzschild black holes). At high energy  $E$ ,

$$\Omega_n \simeq \left[ \frac{cV}{(2\pi)^3} \right]^n \frac{1}{n!} e^{4\pi[E-(n-1)m_0]^2} e^{4\pi(n-1)m_0^2} . \quad (29)$$

Fig. 1. The equilibrium configuration of  $N$  black holes (upper right) is one massive black hole (circled cross) and  $N - 1$  massless black holes. By the bootstrap property, the gas provides a statistical model for a single quantum black hole.

The most probable equilibrium configuration is determined from the condition

$$\left. \frac{d\Omega_n(E, V)}{dn} \right|_{n=N(E, V)} = 0 . \quad (30)$$

The equilibrium state found from this condition is very inhomogeneous. For  $N$  black holes there are one massive and  $N - 1$  massless black holes in the gas (Fig.1). The entropy for such a gas can be approximated by

$$\begin{aligned} S(E, V) &\equiv \ln \Omega(E, V) \simeq \ln \Omega_N(E, V) \\ &\simeq N \ln \left[ \frac{cV}{(2\pi)^3} \right] - \ln \Gamma(N + 1) + S_H(E) , \end{aligned} \quad (31)$$

where  $S_H(E)$  is the Hawking entropy. The microcanonical temperature is

$$\beta = \frac{dS}{dE} = \frac{dS_H}{dE} = 8\pi E . \quad (32)$$

From this expression we see that the microcanonical temperature is the same as the Hawking temperature of the most massive black hole in the ensemble. The microcanonical specific heat turns out to be negative

$$C_V = -\beta^2 \frac{dE}{d\beta} = -\frac{\beta^2}{8\pi} . \quad (33)$$

This does not create a problem, however, because the microcanonical specific heat is allowed in principle to be negative. Our statistical analysis of a gas of Schwarzschild black holes has revealed that it obeys the statistical bootstrap condition

$$\frac{\Omega(E)}{\sigma(E)} \rightarrow 1, \quad E \rightarrow \infty. \quad (34)$$

Such a property is also pictured in Fig. 1. Also we have shown<sup>5</sup> that black hole scattering amplitudes are dual in the sense that the number of open channels grows in parallel with the degeneracy of states as the energy is increased. These two properties are evidence of the conformal nature of black holes as quantum objects.

#### 4. Mean Field Theory

In addition to the WKB approximation there is another semiclassical approximation which can be used to study the properties of black holes. In the mean field approximation fields are quantized on a classical black hole background. Since black holes have a horizon which causally separates two regions of space, we must double the number of degrees of freedom. Two Fock spaces are required. This doubling of the number of degrees of freedom bears a strong resemblance to thermofield dynamics (TFD)<sup>19</sup>.

If we look at the mean field theory, we find that the vacuum for quantum fields scattered off of black holes can be written as

$$|out; 0\rangle = \frac{1}{Z^{1/2}(\beta)} \sum_{n=0}^{\infty} e^{-\beta n \omega / 2} |n\rangle \otimes |\tilde{n}\rangle, \quad (35)$$

where

$$Z = \sum_{n=0}^{\infty} e^{-\beta n \omega}, \quad (36)$$

and the  $|\tilde{n}\rangle$  states provide a basis set for the sector causally disconnected from the observer. Physical operators are defined on the basis set for the states  $|n\rangle$  outside the horizon. The expectation value of a physically observable operator  $\mathcal{O}$  in the *out* region is given as,

$$\langle out; 0 | \mathcal{O} | out; 0 \rangle = \frac{1}{Z(\beta)} \sum_n e^{-n \beta \omega} \langle n | \mathcal{O} | n \rangle, \quad (37)$$

which corresponds to a canonical ensemble average. As before the temperature is determined by the surface gravity<sup>20</sup>

$$\beta = \frac{2\pi}{\kappa} = \beta_H. \quad (38)$$

If the operator  $\mathcal{O}$  is chosen to be the number operator

$$\mathcal{O} = a_k^\dagger a_k , \quad (39)$$

then the expression in Eq.(37) is the number density

$$n_k(m; \beta_H) = \frac{1}{e^{\beta_H \omega_k(m)} - 1} . \quad (40)$$

If black holes are described by a local field theory, this expression presents a problem because it implies loss of coherence. The *in* state is a pure state

$$| in; 0 \rangle = | 0 \rangle \otimes | \tilde{0} \rangle , \quad (41)$$

but the number density obtained from the outgoing states is a thermal distribution.

If black holes are interpreted as quantum excitations of a p-brane, non-local effects must be taken into account. To take account of all possible mass states, we must sum over the mass

$$n_k(\beta_H) = \int_0^\infty dm \, \sigma(m) n_k(m; \beta_H) . \quad (42)$$

The thermal vacuum must now be modified to include contributions from all possible mass states,

$$| out; 0 \rangle = \frac{1}{Z^{1/2}(\beta)} \left[ \prod_{m,k} \sum_{n_{k,m}=0}^\infty \right] \prod_{m,k} e^{-\beta n_{k,m} \omega_{k,m}/2} | n_{k,m} \rangle \otimes | \tilde{n}_{k,m} \rangle , \quad (43)$$

in which the momentum states are shown explicitly. The quantity in square brackets represents the product of sums over the discrete values of the momentum and mass. Changing from discrete to continuous values of the momentum and mass, we find for the canonical partition function

$$Z(\beta) = \exp \left( \frac{-V}{(2\pi)^{D-1}} \int_{-\infty}^\infty d^{D-1} \vec{k} \int_0^\infty dm \, \sigma(m) \ln[1 - e^{-\beta_H \omega_k(m)}] \right) . \quad (44)$$

This expression can be equated to the definition of the partition function to obtain Hagedorn's self-consistency condition<sup>21</sup> for a system of particles in thermodynamical equilibrium

$$\begin{aligned} \int_0^\infty \Omega(E) e^{-\beta E} dE &= \exp \left( \frac{-V}{(2\pi)^{D-1}} \int_{-\infty}^\infty d^{D-1} \vec{k} \right. \\ &\quad \times \left. \int_0^\infty dm \, \sigma(m) \ln[1 - e^{-\beta_H \omega_k(m)}] \right) . \end{aligned} \quad (45)$$

The only objects which satisfy the self-consistency condition are strings<sup>21,22,23</sup>

$$\sigma(m) \sim e^{bm} \quad (m \rightarrow \infty) , \quad (46)$$



for  $\beta_H > b = \text{Hagedorn's inverse temperature}$ . Black holes do not satisfy Hagedorn's condition because

$$\sigma_{BH}(m) \sim e^{C(D)m^{D-2/D-3}}, \quad (47)$$

so that the exponent of  $m$  is always greater than 1 (unless  $D = \infty$ ). The black hole system is not in thermal equilibrium because it is not a self-consistent solution under the assumption of thermal equilibrium. We conclude therefore that the thermal vacuum is the false vacuum for the black hole system.

## 5. Microcanonical Formulation

To determine the true vacuum for a black hole system we begin by writing the thermal vacuum in terms of the density matrix  $\hat{\rho}$

$$|0(\beta)\rangle = \hat{\rho}^{1/2}(\beta, \mathbf{H}) | \mathfrak{S} \rangle, \quad (48)$$

for the case of thermal equilibrium, where

$$\begin{aligned} \hat{\rho}(\beta, \mathbf{H}) &= \frac{\rho(\beta, \mathbf{H})}{\langle \mathfrak{S} | \rho(\beta, \mathbf{H}) | \mathfrak{S} \rangle}, \\ \rho(\beta, \mathbf{H}) &= e^{-\beta \mathbf{H}}, \\ | \mathfrak{S} \rangle &= \left[ \prod_{k,m} \sum_{n_{k,m}=0}^{\infty} \right] \prod_{k,m} |n_{k,m}\rangle \otimes | \tilde{n}_{k,m} \rangle. \end{aligned} \quad (49)$$

The power of  $\frac{1}{2}$  on the density matrix  $\hat{\rho}$  is somewhat arbitrary as will be discussed below.

Observable quantities are obtained by evaluating  $\text{Tr} \mathcal{O}$  where  $\mathcal{O}$  represents any observable operator

$$\text{Tr} \mathcal{O} = \langle \mathfrak{S} | \mathcal{O} | \mathfrak{S} \rangle. \quad (50)$$

In particular we consider the free field propagator, which is defined by the relation

$$\Delta_{\beta,\alpha}^{ab}(x_1, x_2) = -i \langle \mathfrak{S} | T \hat{\rho}^{1-\alpha} \phi^a(x_1) \phi^b(x_2) \hat{\rho}^\alpha | \mathfrak{S} \rangle, \quad (51)$$

where the  $\phi^a$ 's are the so-called thermal doublets

$$\phi^a(x) = \begin{pmatrix} \phi(x) \\ \tilde{\phi}^\dagger(x) \end{pmatrix}, \quad (52)$$

and  $\alpha$  is a parameter which can be chosen between 0 and 1. In writing Eq.(51) use has been made of the invariance of the trace under cyclic rotation of the operators<sup>24</sup>. The Fourier transform of the propagator in Eq.(51) is

$$\Delta_{\beta,\alpha}^{ab} = \frac{\tau_3}{k^2 + m^2 - i\epsilon\tau_3} + \frac{2\pi i \delta(k^2 + m^2)}{e^{\beta|k_0|} - 1} \begin{pmatrix} 1 & e^{(1-\alpha)\beta|k_0|} \\ e^{\alpha\beta|k_0|} & 1 \end{pmatrix}, \quad (53)$$

where  $\tau_3$  is the Pauli matrix

$$\tau_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} . \quad (54)$$

In order to determine the true vacuum for this system it will be convenient to choose  $\alpha = 1$ . In this case the thermal vacuum is given by

$$| \beta \rangle = \hat{\rho}(\beta; \mathbf{H}) | \mathfrak{F} \rangle . \quad (55)$$

We can now formally define the microcanonical vacuum  $| E \rangle$  by the expression

$$| \beta \rangle = \int_0^\infty e^{-\beta E} | E \rangle dE . \quad (56)$$

The microcanonical analysis of the previous sections strongly suggests these considerations. In this basis physical correlation functions are expressed as

$$G_E^{a_1 \dots a_N}(1, 2, \dots, N) = \langle \mathfrak{F} | T \phi^{a_1}(1), \dots, \phi^{a_N}(N) | E \rangle , \quad (57)$$

The normalization for the microcanonical vacuum can be determined from the fact that the thermal vacuum normalization is given by,

$$\langle \mathfrak{F} | \beta \rangle = 1 . \quad (58)$$

Thus according to the definition of  $| E \rangle$  we must have

$$\langle \mathfrak{F} | E \rangle = \delta(E) . \quad (59)$$

An explicit form for the microcanonical vacuum can be formally obtained from the expression for the thermal vacuum with  $\alpha = 1$ , Eq.(55), by an inverse Laplace transform. Using the fact that the partition function obeys the relation

$$Z^{\pm 1}(\beta, V) = \int_0^\infty dE e^{-\beta E} \Omega(E, \pm V) , \quad (60)$$

and the density matrix obeys the relation

$$\rho(\beta, \mathbf{H}) = \int_0^\infty dE e^{-\beta E} \rho(E, \mathbf{H}) , \quad (61)$$

where the microcanonical density matrix is given by

$$\rho(E, \mathbf{H}) = \delta(E - \mathbf{H}) , \quad (62)$$

the microcanonical vacuum  $| E \rangle$  is found to be

$$| E \rangle = \Omega(E - \mathbf{H}, -V) | \mathfrak{F} \rangle , \quad (63)$$

with

$$\begin{aligned}\Omega(E, V) = & \delta(E) + \sum_{n=1}^{\infty} \left( \frac{V}{(2\pi)^{D-1}} \right)^n \frac{1}{n!} \left[ \prod_{i=1}^n \int_0^{\infty} dm_i \sigma(m) \right. \\ & \times \left. \int_{-\infty}^{\infty} d^{D-1} \vec{k}_i \sum_{l_i=1}^{\infty} \frac{1}{l_1, l_2, \dots, l_n} \delta(E - \sum_{i=1}^n l_i \omega_{k_i}(m_i)) \right].\end{aligned}\quad (64)$$

In this expression  $\sigma(m_i)$  is the black hole degeneracy of states and  $\omega_k(m) = \sqrt{k^2 + m^2}$ . The first term in Eq.(64) comes from the vacuum sector ( $E = 0$ ). If  $E > 0$ , then only the second term contributes. We can define a normalized vacuum as

$$|0(E)\rangle = \frac{|E\rangle}{\delta(0)}, \quad (65)$$

so that

$$\langle \mathfrak{S} | 0(E) \rangle = 1, \quad (E = 0). \quad (66)$$

The set of equations (57,59,63,64, and 65) defines our quantum theory of fields in black hole spacetimes. Interaction effects have been neglected up to this point, but they can be taken into account by means of the microcanonical propagator. Using the expression for the 2-point function

$$G_E^{a_1 a_2}(x_1, x_2) = \langle \mathfrak{S} | T \phi^{a_1}(x_1) \phi^{a_2}(x_2) | 0(E) \rangle, \quad (67)$$

and Eq.(66), we can obtain the microcanonical propagator

$$\begin{aligned}\Delta_{E,1}^{ab}(k) = & \frac{1}{\delta(0)} \left( \frac{\tau_3 \delta(E)}{k^2 + m^2 - i\epsilon \tau_3} + 2\pi i \delta(k^2 + m^2) \right. \\ & \times \left[ \sum_{l=1}^{\infty} \delta(E - l|k_0|) \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} + \delta(E) \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right] \end{aligned}\quad (68)$$

Of course only the (1,1)-component  $\Delta_{E,1}^{11}$  is physically observable, and this component is essentially Weldon's propagator<sup>25</sup>.

We can now obtain the particle number density for Hawking radiation for this theory. From Eq(68) we see that

$$n_{k,m} = \sum_l \frac{\delta(E - l\omega_k(m))}{\delta(0)}, \quad (69)$$

The question is: Does this describe a pure state? To answer this question we note that a necessary and sufficient condition for a density matrix to describe a pure state is the idempotency condition

$$\int_0^{\infty} dE' \hat{\rho}_{E_1 E'} \hat{\rho}_{E' E_2} = \hat{\rho}_{E_1 E_2}. \quad (70)$$

In this case

$$\hat{\rho}_{E_1 E_2}(E) = \frac{\delta(E - E_1)\delta(E_1 - E_2)}{\delta(0)} , \quad (71)$$

where

$$\hat{\rho}(E, \mathbf{H}) = \frac{\delta(E - \mathbf{H})}{\delta(0)} , \quad (72)$$

and

$$\int dE \hat{\rho}_{EE} = 1 . \quad (73)$$

Substitution of the expression for  $\hat{\rho}_{E_1 E_2}(E)$  into the idempotency condition shows that this condition is satisfied.

## 6. Conclusions

The foregoing analysis clearly shows that for semiclassical quantization of fields in black hole backgrounds the microcanonical ( $| E >$ ) vacuum is the proper choice, not the thermal vacuum ( $|\beta >$ ). A fixed energy basis for the Hilbert space of the theory should be used instead of the usual thermal state. Black hole states are therefore particle states.

Our analysis also made use of the fact that quantum gravity must be treated as a nonlocal quantum field theory. In our interpretation of black holes as quantum objects the quantum degeneracy of states points to black holes as the excitation modes of  $p$ -branes, with  $p = \frac{D-2}{D-4}$ . The self-consistent treatment of black holes as quantum extended objects implies that black holes are elementary particles.

One interesting extension of the quantum theory of fields in black hole spacetimes given above would be to work out the path integral formulation of field quantization in the fixed energy basis. A further extension is the application of this formalism to string theory. We are also interested in extracting predictions of measurable effects using the thermodoublet formalism of black holes.

## 7. Acknowledgements

This work was supported in part by the Department of Energy under Grant No. DE-FG05-84ER40141.

## 8. References

1. S. Hawking, *Phys. Rev.* **D14** (1976)2460.
2. J.D. Bekenstein, *Phys. Rev.* **D7**(1973)2333; *Phys. Rev.* **D9**(1974)3292.

3. See Ref.[1].
4. J. Preskill, *Proceedings of the International Symposium on Black Holes, Membranes, Wormholes and Superstrings*, The Woodlands, Texas, January, 1992.
5. B. Harms and Y. Leblanc, *Phys. Rev.* **D46**(1992)2334.
6. B. Harms and Y. Leblanc, *Phys. Rev.* **D47**(1993)2438.
7. B. Harms and Y. Leblanc, *Texas/PASCOS 92: Relativistic Astrophysics and Particle Cosmology*, C.W. Ackerlof and M.A. Srednicki eds., Annals of the New York Academy of Sciences **688**(1993)454.
8. B. Harms and Y. Leblanc, to appear in the *Proceedings of SUSY-93*, Northeastern University, Boston, MA (March 1993).
9. B. Harms and Y. Leblanc, University of Alabama preprint no. UAHEP-935.
10. B. Harms and Y. Leblanc, University of Alabama preprint no. UAHEP-936.
11. B. Harms and Y. Leblanc, University of Alabama preprint no. UAHEP-939.
12. J.M. Bardeen, B. Carter and S.W. Hawking, *Comm. Math. Phys.* **31**(1973)161.
13. S.W. Hawking, *Comm. Math. Phys.* **43**(1975)199.
14. G.W. Gibbons and S.W. Hawking, *Phys. Rev.* **D15**(1977)2752.
15. S. Fubini, A.J. Hanson and R. Jackiw, *Phys. Rev.* **D7**(1973)1732.
16. J. Dethlefsen, H.B. Nielsen and H.C. Tze, *Phys. Lett* **48B**(1974)48.
17. A. Strumia and G. Venturi, *Lett. Nuo. Cim.* **13**(1975)337.
18. E. Alvarez and T. Ortin, *Phys. Lett.* **A7**(1992)2889.
19. H. Umezawa, H. Matsumoto and M. Tachiki *Thermo Field Dynamics and Condensed States*, North-Holland Publishing Co. Amsterdam, 1982.
20. N. D. Birrell and P.C.W. Davies, *Quantum Fields in Curved Space*, Cambridge University Press, Cambridge, 1982.
21. R. Hagedorn, *Supp. Nuo. Cim.* **III**(1965)147.
22. S. Frautschi, *Phys. Rev.* **D3**(1971)2821.
23. R.D. Carlitz, *Phys. Rev.* **D5**(1972)3231.
24. T. Arimitsu, H. Umezawa and Y. Yamanaka *J. Math. Phys.* **28**(1987)2741.
25. H.A. Weldon, *Ann. Phys.* **193**(1989)166.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9311077v1>